

notata 28 mm. longa 6.5 mm. diametro subcylindrica parce pilosa sub apice paulo contracta basi dorso semigloboso-inflata, lobis erectis subaequalibus 1.5–2 mm. longis obtusis; antheris omnes inter sese cohaerentibus; bacca globosa apiculata 7 mm. diametro glabra rubra.—JAMAICA: County of Middlesex, St. Ann Parish, trailing over rocks in forest, Mt. Diablo, March 12, 1936, *Hunnewell & Griscom 14481* (TYPE, unicate, in Gray Herb.)

Columnnea Hunnewellii appears to be most closely related to *C. jamaicensis* Urban, from which it differs in its coarsely toothed sepals and much smaller corolla-lobes. Its inclusion in the section *Pterygoloma* Hanst., however, will necessitate the redefinition of that section to include species with dentate sepals.

GRAY HERBARIUM.

BRAYA HUMILIS (C. A. Meyer) Robinson, var. **leiocarpa** (Trautv.), comb. nov. *Sisymbrium nanum* Bunge, var. *leiocarpum* Trautv. in Act. Hort. Petrop. v. 25 (1877). *Pilosella novae-angliae* Rydb. in Torreyia, vii. 158 (1907). *Arabidopsis novae-angliae* (Rydb.) Britton in Britton & Brown, Ill. Fl. ed. 2, ii. 176 (1913), as to type. *B. humilis*, var. *novae-angliae* (Rydb.) Fernald in RHODORA, xx. 202 (1918).—M. L. FERNALD.

Volume 39, no. 462, including pages 193–232, was issued 8 June, 1937.